

MTHE 237 - Lecture Notes

DIFFERENTIAL EQUATIONS FOR ENGINEERING SCIENCE

Prof. Thomas Barthelmé • Fall 2025 • Queen's University

Contents

1	Mathematical modelling & direction field	3
1.1	How to solve	3
1.2	Population Models	3
1.3	Malthusian and competitors	3
1.4	Direction Fields	4
2	Terminology & classification of differential equations	5
2.1	Examples	5
2.2	Types of differential equations	5
2.3	Ordinary differential equations	5
2.4	Confirming a solution of an ODE	5
2.4.1	Superposition principle for linear ODEs	5
3	Initial value problems and the Existence and Uniqueness Theorem for 1st order ODEs	7
3.1	Solving ODEs from implicit relations (template)	7
3.2	Theorem: Existence and Uniqueness of Solutions to the 1st order IVP, or Picard-Lindelof, or Cauchy-Lipshitz	7
4	Initial value problems and the Existence and Uniqueness Theorem for 1st order ODEs (cont.)	9
4.1	Uniqueness fails	9
4.2	Existence fails	9
4.3	Alternate form	9
4.4	Picard Iterations	9
5	Separable Equations	11
5.1	Definitions	11
5.2	Solutions of some ODEs	11
5.3	Separable equations	12
6	Separable Equations (cont.)	13
6.1	Example (implicit solution)	13
6.2	Example (explicit solution)	13
7	Variable Coefficients (Integration Factor Method)	14

7.1	General Solution	14
7.2	Example 1	14
7.3	Example 2	14
8	Exact Equations	16
8.1	Example 1	16
8.2	Example 2	16
8.3	Clairaut's Theorem (Test for exactness)	17
9	Second order linear differential equations	18
9.1	Wronskian	18
9.2	Representation Theorem	18
10	Homogeneous equations with constant coefficients	19
10.1	Case 1: $b^2 - 4ac > 0$	19
10.2	Case 3: $b_2 - 4ac < 0$	19
11	Homogeneous equations with constant coefficients (cont.)	21
11.1	Case 2: $b_2 - 4ac = 0$	21
12	Mechanical and electrical vibrations	22
13	Second order linear nonhomogeneous equations	23
13.1	Particular Solutions	23
13.2	Method of Undetermined Coefficients	23
14	The phenomenon of resonance	25
14.1	Resonance and Generalization	26
14.2	Examples	26
15	Method of variation of parameters	28
16	Making a Particular Solution Guess for Second Order Linear ODEs	29
16.1	Quick Table	29
17	How to Solve an ODE	30
17.1	General Setup	30
17.2	First Order	30
17.2.1	Separation of Variables	30
17.2.2	Variable Coefficients	30
17.2.3	Exact Equations	30
17.2.4	Picard Iteration (Approximate Solutions)	31
17.3	Second Order	31
17.3.1	Characteristic Equation Method (Constant Coefficients)	31
17.3.2	Nonhomogeneous Equations	32

1 Mathematical modelling & direction field

1.1 How to solve

In order to provide a mathematical model describing physical phenomena, we can follow the following steps:

1. Fix our dependent and independent variables and set a frame of references for their measures
2. Choose convenient units of measurements
3. Find the underlying principle governing the motion of the object
4. Rewrite the above relation in terms of the variable we have chosen at step 1
5. Find a solution by integrating both sides
6. Add side conditions to eliminate constants and to find a **unique solution**

Example

$$\begin{aligned}F &= ma \\a &= \frac{dv}{dt} = \frac{d^2h}{dt^2} \\ \frac{d^2h}{dt^2} &= -g \\v(t) &= \frac{dh}{dt} = - \int g dt = -gt + c_1 \\h(t) &= \int v(t) dt = -\frac{1}{2}gt^2 + c_1t + c_2 \\h(0) &= 1, v(0) = 0 \\h(t) &= -\frac{1}{2}gt^2 + 1\end{aligned}$$

1.2 Population Models

Given some internal and external conditions: $\frac{dp}{dt}$ = growth rate - death rate

Death rate = 0

$\frac{dp}{dt} = k_1p$, where $p(0) = p_0$, and $k_1 > 0$ is the proportionality factor for the growth rate

$$\begin{aligned}\frac{1}{p}dp &= k dt \\\ln p &= kt + C \\p(t) &= Ce^{kt}\end{aligned}$$

1.3 Malthusian and competitors

A Malthusian model is a general model for population with rates k_1 and k_2 proportional to p

We can consider more factors to the death rate, called **competitors**, with two-party interactions modelled by $\frac{p(p-1)}{2}$

$$\frac{dp}{dt} = k_1 p - k_3 \frac{p(p-1)}{2}$$

We can rearrange the terms to find an equation in the form of a **logistic model**


$$\frac{dp}{dt} = -\frac{k_3}{2} \left(p^2 - \left(q + \frac{2k_1}{k_3} \right) p \right)$$

which takes the form

$$\frac{dp}{dt} = -Ap(p - p_1), \text{ where } A = \frac{k_3}{2} \text{ and } p_1 = 1 + \frac{2k_1}{k_3}$$

1.4 Direction Fields

For $\frac{dp}{dt} = p(2 - p)$, $\frac{dp}{dt} > 0 \Leftrightarrow 0 < p(t) < 2$



$p = 0, 2$ are equilibria

2 Terminology & classification of differential equations

A differential equation is an equation involving an independent variable (e.g., x , t), a dependent variable, and derivatives of the dependent variables

2.1 Examples

$y' = y + x$, y is dependent and x is independent

$\frac{\partial u}{\partial x} + \frac{\partial u}{\partial y} = x - 2y$, where x, y are independent and the function $u(x, y)$ is dependent

2.2 Types of differential equations

A differential equation with derivatives with respect to one variable is an ODE

A differential equation with partial derivatives with respect to two or more variables is a PDE

2.3 Ordinary differential equations

The general form of an ODE is given by $F(x, y, y', \dots, y^{(n)}) = 0$

The **order** of a differential equation is the order of the highest derivative appearing in the equation

An ODE is **linear** if: 1. y, y', y'', \dots appear only to the first power 2. No product like yy' exist 3. Coefficients $a_i(x)$ can be constants or functions of x , but not of y

A linear ODE of order n has the form: $a_n(x)y^{(n)} + a_{n-1}(x)\frac{d^{n-1}y}{dx^{n-1}} + \dots + a_1x\frac{dy}{dx} + a_0(x)y = f(x)$

2.4 Confirming a solution of an ODE

A function $\phi(x)$ is a solution of an n -th order ODE on some interval $I = (a, b)$ if 1. ϕ is n -times differentiable on I 2. It satisfies the ODE for every $x \in I$

Consider $y'' + \frac{2}{x^2}y = 0$ and the function $\phi(x) = x^2 - \frac{1}{x}$

We can confirm that its derivatives are continuous for all $x \neq 0$ and satisfy the ODE

Thus, the function ϕ is a solution of the given ODE in $(0, \infty)$

2.4.1 Superposition principle for linear ODEs

Say we have an n -th order linear **homogeneous** ODE $a_n(x)y^{(n)} + a_{n-1}(x)\frac{d^{n-1}y}{dx^{n-1}} + \dots + a_1x\frac{dy}{dx} + a_0(x)y = 0$ and suppose $\phi_1(x), \phi_2(x), \dots, \phi_k(x)$ are solutions.

The theorem says any linear combination $\phi(x) = c_1\phi_1(x) + c_2\phi_2(x) + \dots + c_k\phi_k(x)$ is also a solution in I for any choice of arbitrary constants, and the set of solutions is **closed under linear combinations**

These $\phi_k(x)$ solutions, unlike the next example, form a **vector space**, closed under linear combinations

Now, consider a **non-homogeneous** ODE when $f(x) \neq 0$, the linear combination property fails, and only holds in the homogeneous case

3 Initial value problems and the Existence and Uniqueness Theorem for 1st order ODEs

Given the n -th order ODE $F(x, y, y', \dots, y^{(n)}) = 0$ in some interval I , the following two side conditions can be appended to the equation

1. **Initial Conditions:** dependent variable and all its derivatives up to order $n - 1$ are specified at the same point $x_0 \in I$
2. **Boundary Conditions:** only applicable to PDEs

The problem of finding a solution to an ODE in an interval I containing x_0 and *such that the initial conditions are satisfied* is called **initial boundary value problem (IVP)**

3.1 Solving ODEs from implicit relations (template)

Example: given the implicit relation $x_2 + y^2 = 1, x \in (-1, 1)$, we want to see if $y(x)$ is a solution to the ODE $y' = -\frac{x}{y}$

1. **Differentiate w.r.t. dependent variable**

$$2x + 2yy' = 0$$

2. **Solve for y'**

$$y' = -\frac{x}{y}, \text{ this is the ODE}$$

3. **Solve implicit relation explicitly**

$$\text{Rearrange for } y: y(x) = \pm\sqrt{1-x^2}$$

4. **Verify candidates**

For $y(x) = +\sqrt{1-x^2}$, $y' = \frac{-x}{\sqrt{1-x^2}}$, which matches the original ODE For $y(x) = -\sqrt{1-x^2}$, $y' = \frac{x}{\sqrt{1-x^2}}$, which also satisfies the ODE

5. **Apply initial conditions**

$$y_1(x) = -\sqrt{1-x^2} \Rightarrow y_1(0) = -1 \text{ is not valid } y_2(x) = \sqrt{1-x^2} \Rightarrow y_2(0) = 1 \text{ is valid}$$

\therefore the unique solution to the IVP is $y(x) = \sqrt{1-x^2}$


3.2 Theorem: Existence and Uniqueness of Solutions to the 1st order IVP, or Picard-Lindelof, or Cauchy-Lipshitz

Consider the IVP $y' = f(x, y)$ and $y(x_0) = y_0$

If f and $\frac{\partial f}{\partial y}$ are continuous in some rectangle $R = \{(x, y) \in \mathbf{R}^2 : a < x < b, c < y < d\}$, that contains the point (x_0, y_0) , then the IVP admits a unique solutions $\phi : (x_0 - \delta, x_0 + \delta) \rightarrow \mathbf{R}$ for some $\delta > 0$

If $f(x, y)$ is continuous, then at least one solution exists. Continuity of f is enough for *existence*, but not for *uniqueness*

If $f(x, y)$ is continuous and $\frac{\partial f}{\partial y}$ is continuous, then this is enough to show that *uniqueness*



Intuition:

In the original equation, this is like a slope field that tells you the slope of the solution curve at each point (x, y) , $f(x, y)$

If f is continuous, there are no gaps in the slope field. That means you can follow the slope arrows starting at origin to trace out a path

If $\frac{\partial f}{\partial y}$ is also continuous, then nearby paths cannot cross each other. Because the slope field is well defined, you cannot have two different marbles starting at the same point and taking different paths

The rectangle R is the safe zone where f behaves nicely

δ is how far you can move horizontally from x_0 while staying inside the safe zone

Inside $(x_0 - \delta, x_0 + \delta)$, the slope field is well-behaved enough to guarantee a single, smooth solution curve

4 Initial value problems and the Existence and Uniqueness Theorem for 1st order ODEs (cont.)

If one of uniqueness or existence fails, then $\phi(x)$ is not a solution to the ODE

4.1 Uniqueness fails

In $f(x, y) = \frac{y}{x}$, f is undefined at $(0, 0)$, but solutions $y(x) = 0$ and $y(x) = x$ hold and satisfy the IVP with $y(0) = 0$, thus $y(x) = cx$ is a solution for any constant c

This means that there are infinitely many solutions, so **uniqueness fails**

4.2 Existence fails

Now suppose $y(0) = 1$, when $x = 0$ this gives $y(0) = 0$, so **existence fails**

4.3 Alternate form

Given a 1st order linear ODE $a_1(x)y' + a_0(x)y = f(x)$, we can always put it in the form $y'(x) = P(x)y + Q(x)$ by denoting $P(x) = -\frac{a_0(x)}{a_1(x)}$, $Q(x) = \frac{f(x)}{a_1(x)}$

Thus, if $(x_0, y_0) \in \{(x, y) : P(x) \text{ and } Q(x) \text{ are continuous}\}$

Moreover, I can be chosen as the largest interval containing x_0 and such that P, Q are both continuous on I , giving a global, not local, solution

4.4 Picard Iterations

Picard iterations are a method to actually construct the solution that the Existence and Uniqueness Theorem guarantees.

We start at step 0, and iterate integration to construct a sequence of approximations (one per step) that should converge to the unique solution of the IVP

We can rewrite an IVP as an integral equation:

$$y' = f(x, y), y(x) = y_0 + \int_{x_0}^x f(t, y_n(t))dt, \text{ where we don't know } y(t)$$

We can approximate $y(x)$ step by step:

step 0: $\phi_0(x) = y_0$

step 1: $\phi_1(x) = y_0 + \int_{x_0}^x f(s, \phi_0(s)) \, ds$

step 2: $\phi_2(x) = y_0 + \int_{x_0}^x f(s, \phi_1(s)) \, ds$

\vdots

step n: $\phi_n(x) = y_0 + \int_{x_0}^x f(s, \phi_{n-1}(s)) \, ds$

\vdots

If, at some step of the Picard iteration, the new approximation $\phi_{k+1}(x)$ turns out to be exactly the same as the previous one $\phi_k(x)$, then we've already reached the solution

Formally, if there exists $k \in \mathbf{N}$ s.t. $\phi_{k+1}(x) = \phi_k(x)$, then the solution would be given by $\phi_k(x) = y_0 + \int_{x_0}^x f(s, \phi_k(s)) \, ds$

Example:

$$y' = 2t(1 + y), y(0) = 0$$

$$\text{Step 0: } \phi_0(t) = 0$$

$$\text{Step 1: } y'(0) = 2t, \phi_1(t) = \int_0^t 2s \, ds = t^2$$

$$\text{Step 2: } \phi_2(t) = \int_0^t 2s(1 + t^2) \, ds = t^2 + \frac{t^4}{2}$$

$$\text{Step 3: } \phi_3(t) = \int_0^t 2s \left(1 + t^2 + \frac{t^4}{2}\right) \, ds = t^2 + \frac{t^4}{2} + \frac{t^6}{2 \cdot 3}$$

$$\text{Step } n: \phi_n(t) = t^2 + \frac{t^4}{2!} + \frac{t^6}{3!} + \cdots + \frac{t^{2n}}{n!} = \sum_{k=1}^n \frac{t^{2k}}{k!}$$

Hence, the sequence of functions converges if and only if the infinites series given above converges.

We can apply the ratio test to find this:

$$\left| \frac{t^{2(k+1)}}{(k+1)!} \frac{k!}{k+1} \right| \rightarrow 0 \text{ as } k \rightarrow \infty$$

So, the series converges for every t , and $\phi(t) = \sum_{k=1}^{\infty} \frac{t^{2k}}{k!}$

5 Separable Equations

5.1 Definitions

General solution: The family of all possible solutions to the ODE, usually containing arbitrary constants

Explicit solution: A solution where the dependent variable (say y) is written explicitly as a function of the independent variable (say x)

Implicit solution: A solution where the relationship between x and y is given as an equation but not solved explicitly for y

5.2 Solutions of some ODEs

A linear ODE with constant coefficients has the form $y' = ay + b$, where a and b are real constants

We can rewrite this ODE in the equivalent form

$$\frac{1}{y + \frac{b}{a}} \frac{dy}{dt} = a$$

By integrating both sides with respect to t , we get

$$\ln \left| y + \frac{b}{a} \right| = t + C \Leftrightarrow \left| y + \frac{b}{a} \right| = e^C e^{at} \Leftrightarrow y(t) = -\frac{b}{a} \pm c e^{at}, \text{ where } C \text{ and } c \text{ are arbitrary constants}$$

The general solution of an ODE determines an infinite family of curves called **integral curves**, where each curve corresponds to a different value of c . Imposing the initial condition, we are selecting the *unique curve* that passes through the initial point (t_0, y_0)

Example

Consider the classic drag proportional to velocity first-order linear ODE $m \frac{dv}{dt} = mg - \gamma v$

Put in standard form

$$\frac{dv}{dt} = -\frac{\gamma}{m}v + g$$

Rewrite into an equivalent form given

$$\frac{1}{y + \frac{b}{a}} \frac{dy}{dt} = a, \text{ where } a = -\frac{\gamma}{m}, b = g$$

The general solution has the form $v(t) = -\frac{b}{a} + C e^{at}$

Plug back in

$$v(t) = \frac{mg}{\gamma} + C e^{-(\gamma/m)t}, \text{ where } C \text{ is an arbitrary constant}$$

Fix C with an initial condition

$$v(0) = v_0 \quad v_0 = \frac{mg}{\gamma} + C \Rightarrow C = v_0 - \frac{mg}{\gamma}$$

Explicit solution by plugging back C

$$v(t) = \frac{mg}{\gamma} + \left(v_0 - \frac{mg}{\gamma} \right) e^{-\gamma t/m}$$

5.3 Separable equations

We were able to solve the previous ODE because we were able to rewrite it in an equivalent form by **separating the variables**, thus the ODE falls in the class of *separable equations*

Definition: Consider the 1st order ODE $y' = f(x, y)$, if $f(x, y) = g(x)p(y)$, where g and p depends only on x and y , respectively, then the ODE is said to be **separable**

Consider the separable ODE on some interval $I = (a, b)$ $\frac{dy}{dx} = g(x)p(y)$ (*)

We have two cases:

1. If $p(y) \equiv 0$, then $y(x) = c$ constant function on I because the RHS is always 0
2. If $p(y) \neq 0$, then (*) can be rewritten as $\frac{1}{p(y)} \frac{dy}{dx} = g(x)$

Let $H = H(y)$ and $G = G(x)$ be the antiderivatives of $\frac{1}{p(y)}$ and $g(x)$, respectively, from the chain rule we obtain $\frac{dH(y(x))}{dx} = \frac{dH}{dy} \frac{dy}{dx} = \frac{1}{p(y)} \frac{dy}{dx} = \frac{dG}{dx}$

Then, since their derivatives are equal, the functions must differ by a constant: $H(y(x)) = G(x) + c \Leftrightarrow \int \frac{1}{p(y)} dy = \int g(x) dx + c$

This is an implicit formula for the solution to the ODE.

For an explicit solution, we apply the **inverse function** of H to both sides: $H(y) = G(x) + C \Rightarrow H^{-1}(H(y)) = H^{-1}(G(x) + C) \Rightarrow y = H^{-1}(G(x) + C)$

Remark: finding the inverse function depends on H

6 Separable Equations (cont.)

6.1 Example (implicit solution)

Consider $y' = \frac{x}{y \exp(x+2y)}$

It can be rewritten as

$$ye^{2y}y' = xe^{-x}$$

It can then be separated and integrated

$$\int ye^{2y}dy = \int xe^{-x}dy$$

$$\frac{1}{2}ye^{2y} - \frac{1}{4}e^{2y} = -(x+1)e^{-x} \Leftrightarrow e^{2y}(2y-1) = -4e^{-x}(x+1) + c$$

In this case, it is not possible to write y as an explicit function of x . We stop at an **implicit solution**

6.2 Example (explicit solution)

$$y' = (1+y^2)\tan x$$

It can be rewritten as

$$\frac{1}{1+y^2} \frac{dy}{dx} = \tan x$$

It can then be separated and integrated

$$\int \frac{1}{1+y^2} dy = \int \tan x dx \Leftrightarrow \arctan[y(x)] = -\ln|\cos x| + c$$

Integration of $\tan x$

$$\begin{aligned} \int \tan x dx &= \int \frac{\sin x}{\cos x} dx = -\int \frac{1}{u} du \text{ where } u = \cos x \text{ and } du = -\sin x dx \\ &= -\int \frac{1}{u} du = -\ln|u| + C = -\ln|\cos x| + C \end{aligned}$$

7 Variable Coefficients (Integration Factor Method)

Consider a 1st order linear ODE with variable coefficients

$$a_1(x)y' + a_0(x)y = b(x) \Leftrightarrow y' + \frac{a_0(x)}{a_1(x)}y = \frac{b(x)}{a_1(x)} \Leftrightarrow \frac{dy}{dx} + P(x)y = Q(x)$$

If $a_0(x) = a_1'(x)$, then the equation becomes $\frac{d}{dx}(a_1(x))y = b(x)$

Definition: An integrating factor is a special function that we multiply through a first-order linear ODE to make it easier to solve. Its purpose is to turn the left-hand side of the equation into the derivative of a product, so we can integrate directly

7.1 General Solution

Let $\frac{dy}{dx} + P(x)y = Q(x)$

1. We multiply through by an *integrating factor*

$$\mu(x) = e^{\int P(x)dx}$$

2. Multiply the whole equation by $\mu(x)$

$$\mu(x)\frac{dy}{dx} + \mu(x)P(x)y = \mu(x)Q(x) \Leftrightarrow \frac{d}{dx}[\mu(x)y] = \mu(x)Q(x)$$

3. Integrate both sides

$$\mu(x)y = \int \mu(x)Q(x)dx + C$$

4. Solve for $y(x)$

$$y(x) = \frac{1}{\mu(x)} \left(\int \mu(x)Q(x)dx + C \right)$$

General Solution:

$$y(x) = e^{-\int P(x)dx} \left(\int Q(x)e^{\int P(x)dx}dx + C \right)$$

7.2 Example 1

Consider $y' + y = 1$, where $P(x) = Q(x) = 1$

Integrating factor: $e^{\int P(x)dx} = e^x$

Use general solution: $y(x) = e^{-x} \left(\int e^x dx + c \right)$

Solve: $y(x) = 1 + ce^{-x}$

7.3 Example 2

Consider $y' + \frac{3}{x}y = 3x - 2$, $y(1) = 1$, where $P(x) = \frac{3}{x}$, $Q(x) = 3x - 2$

Integrating factor: $\mu(x) = e^{3\ln x} = x^3$

Rewrite ODE: $\frac{d}{dx}[\mu y] = \mu Q \Leftrightarrow \frac{d}{dx}[x^3 y(x)] = 3x^4 - 2x^3$

Integrate on both sides: $x^3 y(x) = \frac{3}{5}x^5 - \frac{2}{4}x^4$

Solve: $y(x) = \frac{3}{5}x^2 - \frac{1}{2}x + \frac{c}{x^3}$

Use initial conditions to find C : $y(1) = 1 = \frac{3}{5} - \frac{1}{2} + C \Leftrightarrow C = \frac{9}{10}$

Final Solution: $y(x) = \frac{3}{5}x^2 - \frac{1}{2}x + \frac{9}{10x^3}$

8 Exact Equations

We start with a first-order ODE: $M(x, y) + N(x, y)\frac{dy}{dx} = 0$

which is equivalent to $M(x, y)dx + N(x, y)dy = 0$

Now, suppose there exists a *potential function* $F(x, y)$ such that $\frac{\partial F}{\partial x} = M(x, y)$, and $\frac{\partial F}{\partial y} = N(x, y)$

Then, if F exists, then the ODE is called exact.

Along any solution curve $y(x)$, if we compute the derivative of $F(x, y(x))$, we get:

$$\frac{d}{dx}F(x, y(x)) = \frac{\partial F}{\partial x} + \frac{\partial F}{\partial y} \frac{dy}{dx}$$

But notice that the right-hand side is exactly the same as $M(x, y) + N(x, y)\frac{dy}{dx}$

Since our ODE says that equals zero, it follows that $\frac{d}{dx}F(x, y(x)) = 0$

That means $F(x, y)$ is constant along solution curves, i.e. $F(x, y) = c$, which is the implicit solution of the ODE

8.1 Example 1

Consider $y' = -\frac{2xy^2+1}{2x^2y}$, where we can rearrange to get $M(x, y) = 2xy^2 - 1$ and $N(x, y) = 2x^2y$

Integrate an equation wrt x :

$$\int (2xy^2 - 1)dx = x^2y^2 - x + g(y)$$

The last term $g(y)$ appears instead of constant c because c can depend on y since we are integrating wrt x

Now, differentiate the latter function wrt y :

$$\frac{\partial F}{\partial y} = 2x^2y + g'(y) = 2x^2y, \text{ where the last term appears from } N(x, y) = 2x^2y$$

We see that $g'(y) = 0$, so $g(y) = c$. Therefore, the given ODE is exact and the general solution is implicitly defined by $x^2y^2 - x = c$ for any arbitrary constant c

8.2 Example 2

Consider $3xy + y^2 + (x^2 + xy)y' = 0$

Integrate $M(x, y)$ wrt x : $F(x, y) = \frac{3}{2}x^2y + xy^2 + g(y)$

Differentiate F wrt y : $\frac{\partial F}{\partial y} = \frac{3}{2}x^2 + 2xy + g'(y) = x^2 + xy$

Rearrange for $g'(y)$: $g'(y) = -\frac{1}{2}x^2 - xy$, which cannot hold because the RHS depends on both variables x and y , while g is only a function of y . Hence, there is no F satisfying for the given ODE, which is then *not exact*

8.3 Clairaut's Theorem (Test for exactness)

An ODE is *exact* in R if and only if $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$ for all $(x, y) \in \mathbf{R}$

Get

$$\begin{cases} \frac{\partial F}{\partial y} = N(x, y) \\ \frac{\partial F}{\partial x} = M(x, y) \end{cases}$$

Since M and N are differentiable wrt x and y , then

$$\begin{cases} \frac{\partial^2 F}{\partial x \partial y} = \frac{\partial N}{\partial x} \\ \frac{\partial^2 F}{\partial y \partial x} = \frac{\partial M}{\partial y} \end{cases}$$

Clairaut's Theorem states that if $F(x, y)$ has continuous second order partial derivatives, then $\frac{\partial^2 F}{\partial x \partial y} = \frac{\partial^2 F}{\partial y \partial x}$

Therefore, $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$ for all $(x, y) \in \mathbf{R}$

9 Second order linear differential equations

CREATE CHEAT SHEET TO SOLVE 1ST ORDER ODES

Let $a_2(x)y'' + a_1(x)y' + a_0(x)y = b(x)$, where the coefficients are functions of x only and are continuous on some open interval $I = (a, b)$

If the ODE = 0, we call it homogeneous.

The superposition principle also applies to second order ODEs.

9.1 Wronskian

A second order ODE needs two initial conditions, and they must not be multiples of one another. Else, solutions y_1, y_2 are linearly dependent on the interval I

We can say that a 2nd order ODE is homogeneous if:

Let y_1, y_2 be solutions on the interval of $a_2(x)y'' + a_1(x)y' + a_0(x)y = 0$. If at some point $x_0 \in (a, b)$ these two solutions satisfy

$$\det \begin{pmatrix} y_1(x_0) & y_2(x_0) \\ y_1'(x_0) & y_2'(x_0) \end{pmatrix} \neq 0$$

and call it **Wronskian of y_1 and y_2 at x_0**

9.2 Representation Theorem

Let $y_1(x), y_2(x)$ be two solutions on the interval (a, b) of $a_2(x)y'' + a_1(x)y' + a_0(x)y = 0$

If at some $x_0 \in (a, b)$, these two solutions satisfy: $W[y_1(x_0), y_2(x_0)] \neq 0$, then $y_1(x)$ and $y_2(x)$ are linearly independent solutions on (a, b)

10 Homogeneous equations with constant coefficients

Consider $ay'' + by' + cy = 0$, where $a, b, c \in \mathbf{R}$

If $\phi(x)$ is a solution to the ODE, then $\phi''(x) = -\frac{b}{a}\phi'(x) - \frac{c}{a}\phi(x)$ for all $x \in \mathbf{R}$

This means that the second derivative is a linear combination of the lower order derivatives

If we have a trial function (educated guess for the solution) $y(x) = e^{rx}$, then we get $y'(x) = re^{rx}$ and $y''(x) = r^2e^{rx}$

Into the ODE, we get $e^{rx}(ar^2 + br + c) = 0$, so e^{rx} is a solution $\Leftrightarrow r$ is a root of the 2nd order polynomial. Thus, r must be a solution to the *characteristic equation* $ar^2 + br + c = 0$

10.1 Case 1: $b^2 - 4ac > 0$

If so, then there exists roots $r_1, r_2 \in \mathbf{R}$ with $r_1 \neq r_2$

r_1 cannot equal r_2 because $W[e^{r_1x}, e^{r_2x}] = (r_2 - r_1)e^{(r_1+r_2)x} \neq 0$

Hence, $y_1(x) = e^{r_1x}$ and $y_2(x) = e^{r_2x}$ are linearly independent solutions, and by the Representation Theorem, $y(x) = c_1e^{r_1x} + c_2e^{r_2x}$, where constants are arbitrary and can be found by imposing initial conditions

Example:

Consider $2y'' + 7y' - 4y = 0$

Characteristic equation: $2r^2 + 7r - 4 = 0$

Confirm determinant > 0 : $r_{1,2} = \frac{-7 \pm \sqrt{49+32}}{4} = -4, \frac{1}{2}$

General solution: $y(x) = c_1e^{-4x} + c_2e^{x/2}$

Impose initial conditions if given

10.2 Case 3: $b^2 - 4ac < 0$

In this case, we have two roots $r_{1,2} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = -\frac{b}{2a} \pm i\frac{\sqrt{4ac - b^2}}{2a} =: \alpha \pm i\beta$

Hence, we find two *complex-valued* linearly independent solutions $y_1, y_2 : \mathbf{R} \rightarrow \mathbf{C}$ $y_1(x) = \exp[(\alpha + i\beta)x]$, $y_2(x) = \exp[(\alpha - i\beta)x]$

We would like to find two *real-valued linearly independent solutions*

Important:

Euler's formula: $e^{i\theta} = \cos \theta + i \sin \theta$

$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}$ and $\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$

Also, if the ODE has real coefficients and the complex-valued function $\phi = u(x) + iv(x)$ is a solution, then real-valued function $u(x) = \operatorname{Re}(\phi(x))$ and complex-valued function $v(x) = \operatorname{Im}(\phi(x))$ are solutions of the same equation

Back to Case 3, from Euler's formula we get: $y_{1,2}(x) = e^{\alpha x}[\cos(\beta x) \pm i \sin(\beta x)]$

Also, be the second remark above, we can conclude that also the functions below are also solutions: $Y_1, Y_2 : \mathbf{R} \rightarrow \mathbf{R}$ $Y_1(x) = \operatorname{Re}(y_1(x)) = \operatorname{Re}(y_2(x)) = e^{\alpha x} \cos(\beta x)$ and $Y_2(x) = \operatorname{Im}(y_1(x)) = -\operatorname{Im}(y_2(x)) = e^{\alpha x} \sin(\beta x)$

$$W[Y_1(x), Y_2(x)] = d$$

$$= e^{\alpha x}$$

Moreover, we can check if these are linearly independent via their Wronskian:

Thus, the solutions above are linearly independent (real-valued) solutions of $ay'' + by' + cy = 0$

Moreover, every other solution takes the form $y(x) = c_1 Y_1(x) + c_2 Y_2(x) = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$

Example:

Consider $y'' - y' + y = 0, y(0) = 1, y'(0) = -2$

Characteristic equation $r^2 - r + 1 = 0$

Solution: $r_{1,2} = \frac{1}{2} \pm i\frac{\sqrt{3}}{2}$

- Linearly independent solutions of the ODE:

$$y_1(x) = e^{x/2} \cos\left(\frac{\sqrt{3}}{2}x\right), \quad y_2(x) = e^{x/2} \sin\left(\frac{\sqrt{3}}{2}x\right).$$

- General solution:

$$y(x) = e^{x/2} \left[c_1 \cos\left(\frac{\sqrt{3}}{2}x\right) + c_2 \sin\left(\frac{\sqrt{3}}{2}x\right) \right].$$

Imposing initial conditions, we get

$$y(x) = e^{x/2} \left[\cos\left(\frac{\sqrt{3}}{2}x\right) - \frac{5\sqrt{3}}{3} \sin\left(\frac{\sqrt{3}}{2}x\right) \right]$$

11 Homogeneous equations with constant coefficients (cont.)

11.1 Case 2: $b_2 - 4ac = 0$

If so, then there exist two roots $r_1, r_2 \in \mathbf{R}$ with $r_1 = r_2$

In this case $c = \frac{b^2}{4a}$ and roots $r_1 = r_2 = -\frac{b}{2a}$. Because we essentially have one root, then the solution $y_1(x) = \exp\left(-\frac{b}{2a}x\right)$ exists

By the Representation Theorem, $y(x) = c_1 y_1(x) + c_2 y_2(x)$. Now that we have $y_1(x)$, how do we find $y_2(x)$ such that $\frac{y_2(x)}{y_1(x)} \neq \text{constant}$

We use the *Method of Reduction of Order*, which works on second order linear homogeneous ODEs with variable coefficients $y'' + p(x)y' + q(x)y = 0$

If $\phi_1(x)$ is a solution to the ODE above, and we want to find $\phi_2(x)$ such that they are linear independent on an interval, we want $\phi_2(x) = g(x)\phi_1(x)$ where $g(x)$ is an unknown function to be found to such that the ODE is satisfied by $\phi_2(x)$

Take $\phi_2(x)$ and take the second derivative: $\phi_2'' = g''\phi_1(x) + 2g'\phi_1'(x) + g\phi_1''(x)$

Replacing in the ODE, we get $\Leftrightarrow \phi_1 v' + (2\phi_1' + p(x)\phi_1)v = 0$, where $v(x) := g'(x)$

We can then solve for $v(x)$ because the corresponding ODE is linear and separable

$$g(x) = \int v(x)dx = \int \frac{\exp\left(-\int p(x)dx\right)}{\phi_1^2(x)} dx$$

Example:

Find a second solution to $x^2 y'' + xy' + \left(x^2 - \frac{1}{4}\right)y = 0$ for $x > 0$ given a solution $y_1(x) = x^{-1/2} \sin x$

Let $y_2(x) = v(x)y_1(x)$, then $y_2'' = v''y_1 + 2v'y_1' + vy_1''$

Goal: find $v(x)$. Replacing the latter in the ODE, we find that

$x^2 y_1 v'' + v'(2x^2 y_1' + xy_1) = 0$ because the y term is the same as the ODE, and it drops out

We can divide further by $x^2 y_1$: $v'' + \left(2\frac{y_1'}{y_1} + \frac{1}{x}\right)v = 0$

Now we need $\frac{y_1'}{y_1}$. Since $y_1'(x) = \frac{\sqrt{x} \cos x - \frac{\sin x}{2\sqrt{x}}}{x}$, then the former is $\frac{\cos x}{\sin x} - \frac{1}{2x}$, which can be simplified to $2\frac{\cos x}{\sin x}$

Subbing back, we get that $v(x)$ has to satisfy the following equation $v'' + 2\frac{\cos x}{\sin x}v' = 0$

Reduce order again: Setting $w(x) := v'(x)$, then $w' = -2\frac{\cos x}{\sin x}w \Rightarrow \frac{w'}{w} = -2\cot x$

Integrate: $\ln |w| = -2\ln(\sin x) + C$, $w = v'(x) = \frac{C_1}{\sin^2 x}$

Integrating to get $v(x)$: $v(x) = -C_1 \cot x + C_2$

Build the solution: $y_2(x) = v(x)y_1(x) = (-C_1 \cot x + C_2)\frac{\sin x}{\sqrt{x}} \Rightarrow y_2(x) = \frac{\cos x}{\sqrt{x}}$

12 Mechanical and electrical vibrations

A **damped mass-spring oscillator** is a physical system constituted by a mass m attached to an elastic spring with stiffness constant k and subject to friction $F_f(t) = -\gamma \frac{dy}{dt}$, where $\gamma \geq 0$ is the damping coefficient

The equation of motion is given by $my'' + \gamma y' + ky = F(t) = 0$ with no external force applied to the body

1. If $\gamma \neq 0$ and $\gamma^2 - 4mk \geq 0$, the system is said to be overdamped

$$r_1, r_2 > 0, y \rightarrow 0, x \rightarrow \infty$$

$$r_1 = 0, r_2 < 0, |y| \rightarrow \infty, x \rightarrow \infty$$

2. If $\gamma \neq 0$ and $\gamma^2 - 4mk < 0$, the system is said to be underdamped

$$\alpha > 0, |y| \rightarrow \infty, y \text{ oscillates to } \infty, x \rightarrow \infty$$

$$\alpha < 0, |y| \rightarrow 0, y \text{ oscillates to } 0, x \rightarrow \infty$$


$$\alpha = 0, y \text{ is periodic, no limit}$$

3. If $\gamma = 0$, the system is said to be undamped

$$r_1 > 0, |y| \rightarrow \infty \text{ exponentially. } x \rightarrow \infty$$

$$r_1 < 0, y \rightarrow 0, x \rightarrow \infty$$

$$r_1 = 0, |y| \rightarrow \infty \text{ linearly. } x \rightarrow \infty$$



13 Second order linear nonhomogeneous equations

Consider the IVP $y'' + p(x)y' + q(x)y = f(x), y(x_0) = Y_0, y'(x_0) = Y_1$

This IVP can be proved with the existence and uniqueness theorem, and an interval $I = (a, b)$ can be found

Furthermore, the superposition principle applies, where solutions ϕ_1 and ϕ_2 forms another solution $\phi(x) = c_1\phi_1 + c_2\phi_2$, which solves for $y'' + p(x)y' + q(x)y = c_1f_1(x) + c_2f_2(x)$

13.1 Particular Solutions

Let $y_c(x) = c_1y_1(x) + c_2y_2(x)$ be a solution to a homogeneous equation, and $y_p(x)$ be a solution of the nonhomogeneous equation

Then, $y(x) = y_c(x) + y_p(x) = c_1y_1(x) + c_2y_2(x) + y_p(x)$

The solution $y_c(x)$ is called the **complementary solution**, whereas the solution $y_p(x)$ is called the **particular solution**

13.2 Method of Undetermined Coefficients

Considering a nonhomogeneous ODE, find $y_p(x)$

Example: $y'' - 3y' - 4y = 3e^{2x}$

Take the homogeneous version and find roots $r_1 = 4, r_2 = -1$

Find the complementary solution by looking for a function of the form $y_p(x) = Ae^{2x}$

Plug in for y , get derivatives, and get $A = -\frac{1}{2}$

General solution: $y(x) = y_c(x) + y_p(x) = c_1e^{4x} + c_2e^{-x} - \frac{1}{2}e^{2x}$

Example: $y'' - 3y' - 4y = 2 \sin x$

We complementary homogeneous solution is $y_c(x) = c_1e^{4x} + c_2e^{-x}$

A solution to the nonhomogeneous solution can be guessed by $y_p(x) = A \cos x + B \sin x$

We always use both sine and cosine terms for a right side like $\sin x$ or $\cos x$ because their derivatives produce each other

We then plug in the derivatives and equate coefficients

$$\begin{cases} \text{cosine: } -5A - 3B = 0 \\ \text{sine: } 3A - 5B - 2 = 0 \end{cases}$$

We can solve for A, B and get a general solution: $y(x) = c_1e^{4x} + c_2e^{-x} + \frac{3}{17} \cos x - \frac{5}{17} \sin x$

Example: $y'' - 3y' - 4y = 4x^2 - 1$

$y_c(x) = c_1e^{4x} + c_2e^{-x}$

For a polynomial right hand side, we try a particular solution of the same order ($Ax^2 + Bx + C$)

We then plug in the derivatives and equate coefficients

$$\begin{cases} \text{x squared:} & 2A - 3B - 4C + 1 = 0 \\ \text{x:} & -6A - 4B = 0 \\ \text{constant:} & A + 1 = 0 \end{cases}$$

14 The phenomenon of resonance

Consider the undamped mass-spring oscillator




Figure 3.7: Forced mass-spring oscillator.

General ODE:

$$y'' + \omega^2 y = F_0 \cos(\gamma t), \quad \omega^2 = \frac{k}{m}$$

- ω : Natural frequency (system's own oscillation)
- γ : Forcing frequency (external input)

General Solution (Non-Resonant):

$$y(t) = c_1 \cos(\omega t) + c_2 \sin(\omega t) + y_p(t)$$

Try particular solution:

$$y_p(t) = A \cos(\gamma t) + B \sin(\gamma t)$$

After differentiating and substituting:

$$A(\omega^2 - \gamma^2) \cos(\gamma t) + B(\omega^2 - \gamma^2) \sin(\gamma t) = F_0 \cos(\gamma t)$$

So:

$$A = \frac{F_0}{\omega^2 - \gamma^2}, \quad B = 0$$

Final non-resonant solution (for $\omega \neq \gamma$):

$$y(t) = c_1 \cos(\omega t) + c_2 \sin(\omega t) + \frac{F_0}{\omega^2 - \gamma^2} \cos(\gamma t)$$

14.1 Resonance and Generalization

Resonance occurs when $\omega = \gamma$, i.e., the forcing matches the system's own frequency. This breaks the usual method since the input matches a homogeneous solution.

- **General principle:** When the forcing function is also a solution of the homogeneous equation, *multiply your usual particular solution guess by t for linear independence.*
- For resonance ($\omega = \gamma$), trial:

$$y_p(t) = [A \cos(\omega t) + B \sin(\omega t)]t$$

Solving gives:

$$A = 0, \quad B = \frac{F_0}{2\omega}$$

So the resonance particular solution is:

$$y_p(t) = \frac{F_0}{2\omega} t \sin(\omega t)$$

$f(x)$	$y_p(x)$
$P_n(x) := a_n x^n + \cdots + a_1 x + a_0$	$x^s [A_n x^n + A_{n-1} x^{n-1} + \cdots + A_1 x + A_0]$ Three possibilities for s : $s = 0$ if $r = 0$ is not a solution of the characteristic equation $s = 1$ if $r = 0$ is only one solution of the characteristic equation $s = 2$ if $r = 0$ is double solution of the characteristic equation
$P_n(x)e^{\alpha x}$	$x^s [A_n x^n + A_{n-1} x^{n-1} + \cdots + A_1 x + A_0]e^{\alpha x}$ Three possibilities for s : $s = 0$ if $r = \alpha$ is not a solution of the characteristic equation $s = 1$ if $r = \alpha$ is only one solution of the characteristic equation $s = 2$ if $r = \alpha$ is double solution of the characteristic equation
$P_n(x)e^{\alpha x} \cos(\beta x) + P_m(x)e^{\alpha x} \sin(\beta x)$	$x^s [A_k x^k + A_{k-1} x^{k-1} + \cdots + A_1 x + A_0]e^{\alpha x} \cos(\beta x) +$ $+ x^s [B_k x^k + B_{k-1} x^{k-1} + \cdots + B_1 x + B_0]e^{\alpha x} \sin(\beta x)$ $k = \max\{n, m\}$ Two possibilities for s : $s = 0$ if $r = \alpha + i\beta$ is not a solution of the characteristic equation $s = 1$ if $r = \alpha + i\beta$ is a solution of the characteristic equation

14.2 Examples

Example: $y'' - 3y' - 4y = -8e^{-x}$

$$y_c(x) = c_1 e^{4x} + c_2 e^{-x}$$

Notice the forcing term $f(x) = -8e^{-x}$ is a constant multiple of $c_2 e^{-x}$, thus we need a particular solution of the form Axe^{-x}

We then plug in the derivatives and get the general solution $y(x) = c_1 e^{4x} + c_2 e^{-x} + \frac{8}{5} x e^{-x}$

Example: $y'' - 2y' + y = (x + 1)e^x$

$$y_c(x) = (c_1 + c_2 x)e^x$$

If we plug in the RHS, the DE will equal 0

Then, we look for a particular solution of the form $y_p(x) = x^2(Ax + B)e^x$

Example: $2y'' + 3y' + y = x^2 + 3\sin x$

In this case, we can develop a particular solution from the superposition principle because the forcing term is an addition: $f(x) = x^2 + 3\sin x$

Thus $y_p(x) = y_{p,1}(x) + y_{p,2}(x)$, where the former is a particular solution to $ODE = x^2$ and the latter is a particular solution to $ODE = 3\sin x$

15 Method of variation of parameters

Find a solution to $y'' + 4y = 3 \csc x$, but $f(x)$ cannot be expressed as an l.c. of our common terms from the previous note

We can use **variation of parameters**, which works for any possible type of forcing term

Consider general form ODE $y'' + a(x)y' + b(x)y = f(x)$, and consider the particular solution of the form $y_p(x) = v_1(x)y_1(x) + v_2(x)y_2(x)$

Get $y'_p = v'_1y_1 + v_1y'_1 + v'_2y_2 + v_2y'_2$, and assume that $v'_1y_1 + v'_2y_2 = 0$

Then, $y''_p = v'_1y'_1 + v_1y''_1 + v'_2y'_2 + v_2y''_2$

We can impose that to the ODE and get that $y_p = v_1(x)y_1(x) + v_2(x)y_2(x)$ is a particular solution iff v_1 and v_2 satisfy:

$$\begin{cases} v'_1y_1 + v'_2y_2 = 0 \\ v'_1y'_1 + v'_2y'_2 = f(x) \end{cases}$$

where $W[y_1(x), y_2(x)] \neq 0$

We then find the following solution of the system: $v_1(x) = \int \frac{-f(x)y_2(x)}{W[y_1(x), y_2(x)]} dx$ $v_2(x) = \int \frac{f(x)y_1(x)}{W[y_1(x), y_2(x)]} dx$

16 Making a Particular Solution Guess for Second Order Linear ODEs

Step 1: Write the nonhomogeneous ODE in standard form.

- E.g., $y'' + ay' + by = f(x)$

Step 2: Identify the type of forcing (right-hand side) $f(x)$:

- **Polynomial** ($ax^n + \dots$): Guess another polynomial of the same degree.
- **Exponential** (Ae^{kx}): Guess $y_p = Ae^{kx}$.
- **Trig (sin/cos)** ($A \sin(kx), B \cos(kx)$): Guess $y_p = A \cos(kx) + B \sin(kx)$.
- **Products**: Multiply the corresponding guesses, e.g. xe^{kx} leads to $(Ax + B)e^{kx}$.

Step 3: Check for overlap with homogeneous solutions (roots of characteristic equation):

- Find roots (frequencies, exponentials) of the homogeneous equation.
- If your guess matches a homogeneous solution (same root/frequency):
 - **Multiply your guess by x** (or t if variable is time) for linear independence.
 - If root has multiplicity n , multiply by x^n .

Step 4: Plug your guess into the ODE, solve for coefficients.

- Substitute, collect terms, build a system for unknowns.

16.1 Quick Table

Forcing Function	Standard Guess	If matches homogeneous?
Ax^n	Polynomial of degree n	Multiply by x or higher
Ae^{kx}	Ae^{kx}	Multiply by x or higher
$A \sin(kx), B \cos(kx)$	$A \sin(kx) + B \cos(kx)$	Multiply by x or higher
$x^r e^{kx}$	$(Ax^r + \dots)e^{kx}$	Multiply by x, x^2

Mnemonic:

- If your guess is wiped out by the homogeneous equation, multiply by x for new solutions

17 How to Solve an ODE

17.1 General Setup

ODE form: $F(x, y, y', y'', \dots, y^{(n)}) = 0$

General solution: Family of solutions with arbitrary constants.

Particular solution: Obtained by applying initial/boundary conditions.

Explicit solution: $y = f(x)$.

Implicit solution: Relation between x and y , not fully solved for y .

17.2 First Order

17.2.1 Separation of Variables

Form: $\frac{dy}{dx} = g(x)p(y)$

Steps:

1. Rearrange: $\frac{1}{p(y)}dy = g(x)dx$
2. Integrate both sides
3. Get implicit solution $H(y) = G(x) + C$
4. Solve for y if possible (explicit solution)

17.2.2 Variable Coefficients

Form: $y' = P(x)y = Q(x)$

Steps:

1. Compute integrating factor $\mu(x) = e^{\int P(x)dx}$
2. Multiply ODE by $\mu(x)$: $\frac{d}{dx}[\mu(x)y] = \mu(x)Q(x)$
3. Integrate both sides: $\mu(x)y = \int \mu(x)Q(x)dx + C$
4. Solve for $y(x)$

17.2.3 Exact Equations

Form: $M(x, y) + N(x, y)y' = 0$ or $M(x, y)dx + N(x, y)dy = 0$

Condition (Clairaut's Test): $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$

Steps:

1. Check exactness with Clairaut's test
2. Find potential function $F(x, y)$ such that $\frac{\partial F}{\partial x} = M$, $\frac{\partial F}{\partial y} = N$
3. Implicit solution: $F(x, y) = C$

17.2.4 Picard Iteration (Approximate Solutions)

Form: For IVP $y' = f(x, y), y(x_0) = 0$

Steps:

- Rewrite as integral equation $y(x) = y_0 + \int_{x_0}^x f(t, y(t))dt$
- Approximate iteratively $\phi_{n+1}(x) = y_0 + \int_{x_0}^x f(t, \phi_n(t))dt$
- Sequence converges to the solution

17.3 Second Order

We usually deal with **linear, constant-coefficient ODEs** of the form:

$$ay'' + by' + cy = 0 \quad \text{or more generally} \quad y'' + P(x)y' + Q(x)y = 0$$

17.3.1 Characteristic Equation Method (Constant Coefficients)

1. Write characteristic polynomial:

$$ar^2 + br + c = 0$$

2. Compute discriminant:

$$\Delta = b^2 - 4ac$$

3. Solve according to cases:

Case 1: Two distinct real roots $\Delta > 0$

- Roots:

$$r_{1,2} = \frac{-b \pm \sqrt{\Delta}}{2a}$$

- General solution:

$$y(x) = c_1 e^{r_1 x} + c_2 e^{r_2 x}$$

Case 2: Repeated root $\Delta = 0$

- Root:

$$r = -\frac{b}{2a}$$

- General solution:

$$y(x) = (c_1 + c_2 x)e^{rx}$$

- (Alternate derivation: reduction of order — given one solution y_1 , the second is

$$y_2 = y_1(x) \int \frac{\exp(-\int P(x)dx)}{y_1(x)^2} dx$$

for ODEs in the form $(y'' + P(x)y' + Q(x)y = 0)$)

Case 3: Complex conjugate roots $\Delta < 0$

- Roots:

$$r_{1,2} = \alpha \pm i\beta, \quad \alpha = -\frac{b}{2a}, \quad \beta = \frac{\sqrt{4ac-b^2}}{2a}$$

- General solution:

$$y(x) = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

17.3.2 Nonhomogeneous Equations

For $ay'' + by' + cy = f(x)$:

- General solution = **Complementary solution** (y_c) + **Particular solution** (y_p).

1. Solve homogeneous equation \rightarrow get y_c .
2. Guess form of y_p (method of undetermined coefficients) or use **variation of parameters**.

- Variation of parameters formula for $y'' + P(x)y' + Q(x)y = R(x)$:

$$y_p = u_1(x)y_1(x) + u_2(x)y_2(x)$$

where

$$u'_1 = -\frac{y_2(x)R(x)}{W(y_1, y_2)}, \quad u'_2 = \frac{y_1(x)R(x)}{W(y_1, y_2)}$$

$$\text{and } W(y_1, y_2) = y_1 y'_2 - y'_1 y_2$$